

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)**ScienceDirect**journal homepage: [www.elsevier.com/locate/ajps](http://www.elsevier.com/locate/ajps)**P-071**

## Synthesis, characterization, antibacterial and anticancer evaluation of some novel flavone-3-ols

*Aravinda Pai* \*, *D. Vijay Kumar*, *B.S. Jayashree*

Department of Pharmaceutical Chemistry, Manipal College of Pharmaceutical Sciences, Manipal, Karnataka 576104, India

### ARTICLE INFO

#### Article history:

Available online 26 November 2015

#### Keywords:

Flavonols

Antibacterial

Anticancer

HeLa

V79

A series of novel flavonols were synthesized by cyclizing corresponding hydroxy chalcones using Algar flynn Oyamada reaction [1]. The synthesized compounds were characterized by various spectrochemical methods including IR, MASS, NMR spectroscopy. Out of 13 compounds screened for antibacterial activity [2], compound VMF 12 (a bromo derivative) showed activity with an  $IC_{50}$  value at  $0.16 \mu M$  against *S. aureus*. Compounds VMF 1 and VMF 2 (dichloro derivatives) showed activity with  $IC_{50}$  at  $0.34$  and  $0.46 \mu M$  respectively against *E. coli*. Compound VMF 8 (a dichloro derivative of phenyl substituted  $\gamma$ -benzopyrone) exhibited maximum activity with an  $IC_{50}$  value at  $0.02 \mu M$  concentration against *P. aeruginosa*. None of the compounds were active against *B. subtilis*.

The anti-cancer potency of the test compounds were evaluated by MTT [3] assay for 13 test compounds on two different cell line such as HeLa and V79 cell lines. Compounds such as VMF 7(dichloro derivative) and VMF 8(dichloro derivative) showed  $IC_{50}$  at  $0.38 \mu M$  each, compound VMF 4 (a dichloro derivative)

showed  $IC_{50}$  at  $0.43 \mu M$  against standard Cisplatin and Curcumin, showing an  $IC_{50}$  of  $0.01$  and  $0.12 \mu M$  respectively against HeLa cell lines.

The cytotoxic activity for the test compounds were also performed against V79 cell lines using MTT assay method. Out of the 13 compounds screened, compounds VMF 4 (a dichloro derivative), VMF 9 (a dichloro derivative), VMF 10 (a di bromo derivative), and VMF 13 (a di bromo derivative) showed moderate toxicity with an  $IC_{50}$  greater than  $1 \mu M$ , whereas, other test compounds showed  $IC_{50}$  greater than  $2 \mu M$  against standard Cisplatin and Curcumin with their  $IC_{50}$  value at  $0.15$  and  $0.19 \mu M$  respectively against V79 cell lines.

### Acknowledgement

The authors acknowledge the support received by Manipal University to attend the conference.

\* E-mail address: [aravind.pai@manipal.edu](mailto:aravind.pai@manipal.edu).

Peer review under responsibility of Shenyang Pharmaceutical University.

<http://dx.doi.org/10.1016/j.ajps.2015.11.044>

1818-0876/© 2016 Production and hosting by Elsevier B.V. on behalf of Shenyang Pharmaceutical University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – (A) IR spectra of compound VMF 9, and (B) NMR spectra of compound VMF 3.

## REFERENCES

- [1] Algar J, Flynn JP. Proc Roy Irish Acad 1934;42B:I.
- [2] Oya BD, Meltem C, Nurten A, et al. Synthesis and antimicrobial activities of some new flavone derivatives. Acta Pharmaceutica Turcica 2003;45(1): 31-35.
- [3] Wall NR, O'Connor DS, Plescia J, et al. Suppression of survivin phosphorylation on Thr<sup>34</sup> by flavopiridol enhances tumor cell apoptosis. Cancer Res 2003;63:230-235.